



HEAT DETECTION

Select Synch




Select Synch + CIDR®



PG 6-day CIDR®


Heat detect and AI days 0 to 3. Administer CIDR to non-responders and heat detect and AI days 9 to 12. Protocol may be used in heifers.



HEAT DETECT & TIME AI (TAI)


Select Synch & TAI

Heat detect and AI day 6 to 10 and TAI all non-responders 72 - 84 hr after PG with GnRH at TAI.




Select Synch + CIDR® & TAI

Heat detect and AI day 7 to 10 and TAI all non-responders 72 - 84 hr after PG with GnRH at TAI.



PG 6-day CIDR® & TAI


Heat detect & AI days 0 to 3. Administer CIDR to non-responders & heat detect and AI days 9 to 12. TAI non-responders 72 - 84 hr after CIDR removal with GnRH at AI. Protocol may be used in heifers.



FIXED-TIME AI (TAI)*


7-day CO-Synch + CIDR®

Perform TAI at 60 to 66 hr after PG with GnRH at TAI.



5-day CO-Synch + CIDR®

Perform TAI at 72 ± 2 hr after CIDR removal with GnRH at TAI. Two injections of PG 8 ± 2 hr apart are required for this protocol.




FIXED-TIME AI (TAI)*

for *Bos Indicus* cows only

Bos Indicus PG 5-day + CIDR®


Perform TAI at 66 ± 2 hr after CIDR removal with GnRH at TAI.




* The time listed for "Fixed-time AI" should be considered as the approximate average time of insemination. This should be based on the number of cows to inseminate, labor, and facilities.

HEAT DETECTION


1 Shot PG



7-day CIDR®-PG




MGA®-PG



HEAT DETECT & TIME AI (TAI)


Select Synch + CIDR® & TAI

Heat detect and AI day 7 to 10 and TAI all non-responders 72 - 84 hr after PG with GnRH at TAI.




MGA®-PG & TAI

Heat detect and AI day 33 to 36 and TAI all non-responders 72 - 84 hrs after PG with GnRH at TAI.



14-day CIDR®-PG & TAI

Heat detect and AI day 30 to 33 and TAI all non-responders 72 hrs after PG with GnRH at TAI.




FIXED-TIME AI (TAI)*

Short-term Protocols


7-day CO-Synch + CIDR®

Perform TAI at 54 ± 2 hr after PG with GnRH at TAI.



5-day CO-Synch + CIDR®


Perform TAI at 60 ± 4 hr after CIDR removal with GnRH at TAI. Two injections of PG 8 ± 2 hr apart are required for this protocol.



Long-term Protocols


14-day CIDR®-PG

Perform TAI at 66 ± 2 hr after PG with GnRH at TAI.



MGA®-PG

Perform TAI at 72 ± 2 hr after PG with GnRH at TAI.



* The times listed for "Fixed-time AI" should be considered as the approximate average time of insemination. This should be based on the number of heifers to inseminate, labor, and facilities.

These protocol sheets were assembled by the *Beef Reproduction Task Force*; *BeefRepro.org*. Programs are intended to promote sustainable food production systems by the beef industry through sound reproductive management practices for use in replacement heifers and postpartum cows. The Beef Reproduction Task Force recommends working with a licensed veterinarian for proper use and application of all reproductive hormones. **Approved 11-17-2020.**

GnRH Cystorelin®, Factrel®, Fertagyl®, OvaCyst®, GONABreed®

PG estroPLAN®, Estrumate®, In-Synch®, Lutalyse®, Lutalyse® HighCon, ProstaMate®, SYNCHSURE™